

Results from Searching for Muonium to Antimuonium Conversion

K. JUNGSMANN, A. GROSSMANN, J. MERKEL, V. MEYER, G. ZU PUTLITZ,
I. REINHARD, K. TRÄGER, P.V. SCHMIDT, L. WILLMANN
Physics Institute, University of Heidelberg, D-69120 Heidelberg

R. ENGFER, H.P. WIRTZ
Physics Institute, University of Zürich, CH-8057

R. ABELA, W. BERTL, D. RENKER, H.K. WALTER
Paul Scherrer Institute; CH-5232 Villigen

V. KARPUCHIN, I. KISEL, A. KORENCHENKO, S. KORENCHENKO,
N. KRAVCHUK, N. KUCHINSKY, A. MOISEENKO
JINR Dubna, RU-141980 Dubna

J. BAGATURIA, D. MZAVIA, T. SAKHELASHVILI
Tbilisi State University, GUS-380086 Tbilisi

V.W. HUGHES
Yale University, New Haven, CT 06520

A new result from searching for muonium to antimuonium conversion is reported which sets an upper limit on the coupling constant in an assumed (V-A)×(V-A) type interaction of $G_{M\overline{M}} \leq 3 \cdot 10^{-3} G_F$ (90%C.L.). A particular Z_8 and a minimal 331 GUT model can be ruled out. Further new and stringent limits can be set for masses of bileptonic gauge bosons and λ parameters in R-parity breaking supersymmetric models.

Due to the close confinement in the bound state the hydrogen-like muonium atom ($M = \mu^+ e^-$) is a well suited system to test the fundamental interactions between two leptons from different generations. Of particular interest are searches for lepton number violating processes which occur naturally in many speculative extensions to the standard model. A spontaneous conversion of muonium to antimuonium ($\overline{M} = \mu^- e^+$) would violate additive lepton number conservation by two units and is of interest particularly to e.g. left-right symmetric^{1,2}, supersymmetric^{3,4} and GUT^{5,6,7} models.

At the Paul Scherrer Institute (PSI) in Villigen, Switzerland, an experiment has been successfully completed very recently which searched sensitively for $M - \overline{M}$ conversion. The coincident detection of both of the constituents (μ^- and e^+) of the antiatom in its decay was required as a signature for the conversion process⁸.

A μ^- -decay ($\mu^- \rightarrow e^- + \bar{\nu}_e + \nu_\mu$) releases an energetic electron with an energy spectrum ranging up to 53 MeV and can be identified in a magnetic spectrometer consisting of five cylindrical proportional wire chambers in coaxial geometry operated at 0.1 T field. The positron in the atomic shell of the antiatom is left behind after the decay with an average kinetic energy of $R_\mu=13.5$ eV. This particle is accelerated to 7 keV and transported in a magnetic guiding field onto a position sensitive microchannel plate detector (MCP). The 511 keV radiation from the annihilation of positrons in the MCP can be detected in a segmented pure CsI crystal detector.

The muonium atoms are formed by stopping a beam of positive muons in a SiO_2 powder. A fraction of about 7% of the incoming muons is converted into muonium atoms which diffuse to the surface and leave the powder with thermal energies into the surrounding vacuum. It is essential for the experiment to have the atoms in vacuum, since the $M-\bar{M}$ -oscillation is strongly suppressed in gases or condensed matter due to the removal of symmetry between the atom and the antiatom. The muonium yield has been monitored regularly by reversing the electric acceleration field and all magnetic fields approximately every five hours and by observing energetic positrons and electrons respectively. These measurements also serve for calibrating detector subsystems. The targets are replaced about every third day to account for an observed time dependent decrease of the muonium production efficiency, due to target surface structure and contaminations. Attention has been paid to keep the apparatus as much as possible symmetric for detecting muonium and antimuonium decays.

In the analysis of the final data taking period, which included 1290h of apparatus lifetime, one event was found within 3 standard deviations of all relevant parameters. This is in good agreement with an expected background of 1.7(2) events from accidental coincidences. The physical background due to the allowed process $\mu^- \rightarrow e^- + \bar{\nu}_e + \nu_\mu + e^+ + e^-$ which could fake an event is fully suppressed by a proper choice of the magnetic positron transport system parameters which includes a 90° bend. The combination of all measurement periods encompassing some 1600 h measurement time yields an upper limit on the conversion probability of $P_{M\bar{M}} \leq 8.0 \cdot 10^{-11}/S_b$ (90% C.L.), where S_b reflects the magnetic field dependence of the process and is for 0.1 T in this experiment 0.35 for $(V\pm A)\times(V\pm A)$ and 0.75 for $(V\mp A)\times(V\pm A)$ interaction types¹⁰. This constitutes an improvement by a factor of 2500 over the value established in preceding experiments⁹. This corresponds for an assumed effective four fermion in $(V-A)\times(V-A)$ interaction to a limit on the coupling constant of $G_{M\bar{M}} \leq 3 \cdot 10^{-3} G_F$, with G_F being the weak interaction Fermi coupling constant.

This result rules out a Z_8 model with radiative mass generation by heavy lepton seed⁶. It allows further to set a lower limit on the mass of bileptonic gauge bosons in GUT theories of $2.6 \text{ TeV}/c^2 * g_{3l}$, where g_{3l} depends on the particular symmetry and is of order unity; this is significantly above values extracted from high energy Bhabha scattering experiments. In a minimal 331 model the mass limit on the bilepton an $800 \text{ GeV}/c^2$ upper mass bound has been found. The new limit from this $M\overline{M}$ experiment corresponds to a $850 \text{ GeV}/c^2$ lower bound and excludes the minimal version of this theory. However, the model would still be viable, if one add an octet of Higgs¹².

In the framework of R-parity violating supersymmetric models^{3,4} the bound on the coupling parameters could be lowered by a factor of 15 to $|\lambda_{132}\lambda_{231}| \leq 3 * 10^{-4}$ with assumed superpartner masses of order $100 \text{ GeV}/c^2$ ¹¹. The achieved level of sensitivity allows to narrow slightly the interval of allowed heavy muon neutrino masses in minimal left-right symmetric models¹ (which have predicted a lower bound on $G_{\overline{M}M}$ provided muon neutrinos have a mass above 35 keV) to $\approx 40 \text{ keV}/c^2$ up to the present experimental bound at $170 \text{ keV}/c^2$.

The work is supported in part by the BMBF of Germany, the Russian RFFR and the Schweizer Nationalfond and a NATO research grant.

References

1. P. Herczeg and R.N. Mohapatra, *Phys. Rev. Lett.* **69**, 2475 (1992).
2. A. Halprin, *Phys. Rev. Lett.* **48**, 1313 (1982).
3. R.N. Mohapatra, *Z. Phys. C* **56**, 117 (1992).
4. A. Halprin and A. Masiero, *Phys. Rev. D* **48**, 2987 (1993).
5. H. Fujii et al., *Phys. Rev. D* **49**, 559 (1994).
6. W.S. Hou and G.C. Wong, *Phys. Rev. D* **53**, 1527 (1996).
7. P. Frampton, hep-ph/9711281 (1997) and references therein.
8. R. Abela et al., *Phys. Rev. Lett.* **77**, 1950 (1996).
9. B.E. Matthias et al., *Phys. Rev. Lett.* **66**, 2716 (1991); see also: V.A. Gordeev et al., *JETP Lett.* **59**, 589 (1994).
10. G.G. Wong and W.S. Hou, *Phys. Lett. B* **357**, 145 (1995); K. Horikawa and K. Sasaki, *Phys. Rev. D* **53**, 560 (1996).
11. R. Mohapatra, private communication (1998).
12. P. Frampton, private communication (1998).